SMALL-SCALE OCEAN ACOUSTIC TOMOGRAPHY

Jerald W. Caruthers Naval Research Laboratory Stennis Space Center, MS 39529

Phone: (228)688-5438 fax: (228)688-5341 email: jwc@nrlssc.navy.mil Award # N0001497WX40088

LONG-TERM GOALS

The goal of this project is to establish novel techniques in Ocean Acoustic Tomography.

OBJECTIVES

The objectives are to investigate and extend techniques in OAT based on evolving approaches pursued by Russian investigators and apply them to small-scale tomography of bubble plumes.

APPROACH

Russian scientists have taken a broader view of Ocean Acoustic Tomography (OAT) than has US scientists. Their view equates OAT with essentially any acoustical oceanography inverse problem based on a solution of a system of equations involving any set of acoustic measurements. US work has concentrated more on traditional travel-time and attenuation tomography. In our review of OAT, we have identified several new approaches that would be useful to pursue. We are exploring three applications to small-scale reconstructions involving bubble clouds in shallow water: emission tomography of bubble noise, diffraction tomography for bubble clouds, and non-reciprocity tomography for small-scale motion. The first two are conducted as a joint effort with Dr. Alexander I. Khilko of the Institute of Applied Physics, RAS (Russian Academy of Sciences) and the latter one with Dr. Oleg Godin of the Oceanography Institute, RAS.

The applications of tomography to small-scale reconstruction of bubble cloud properties are being studied in the context of a high-frequency acoustic, shallow-water bubble experiment conducted off the Scripps Pier in San Diego in March of 1997. Data for these efforts were taken using a device called the Delta Frame'that measured acoustic properties at multiple frequencies over 16 paths within a 10-m equilateral triangle.

WORK COMPLETED

Application of standard (attenuation) tomography to reconstructing the bubble clouds has already been completed. A model has been developed for the application of emission tomography to the reconstruction of small-scale noise sources represented by bubble plumes formed just below the surface due to breaking waves. Several papers were presented on this topic. Diffraction tomographic algorithms have been developed. Numerous papers in each of the areas have been written and a draft book has been prepared. The book should go to press early next year.

Several approaches have been developed to extract the current-related information from travel times along not quite reciprocal paths in the Delta Frame experiment geometry. The

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to completing and reviewing the collect this burden, to Washington Headquuld be aware that notwithstanding and DMB control number.	tion of information. Send commen larters Services, Directorate for In:	ts regarding this burden estimate formation Operations and Reports	or any other aspect of to s, 1215 Jefferson Davis	his collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Small-scale Ocean Acoustic Tomography				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Stennis Space Center, MS, 39529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release; distribut	ion unlimited			
13. SUPPLEMENTARY NO	OTES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC	CATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	3	RESPONSIBLE PERSON

Report Documentation Page

Form Approved OMB No. 0704-0188 propagation paths are orders of magnitude shorter and the environment is much more volatile than in other reciprocal transmission experiments. Therefore, *a priori* there is a considerable uncertainty in what to expect from the data on non-reciprocity, particularly from not quite reciprocal paths. The approaches differ by the way effects of variations in sound speed are suppressed and (partially) separated from those due to fluid motion and are based on different assumptions about space- and time-scales of variations of the sound speed and density in the experiment.

RESULTS

Attenuation data from the Delta Frame have been processed and successfully applied in standard tomography. For the more elaborate techniques, data processing is not yet complete.

IMPACT/APPLICATIONS

New techniques in OAT have direct application to sensing the oceanic environments for numerous military and civilian applications.

TRANSITIONS

Techniques in emission, diffraction, and non-reciprocity tomography applied at the small scale are transitioning to applications of sensing bubble clouds in shallow water. Application of such work should be useful in MCM and special warfare.

RELATED PROJECTS

This project is jointly funded by the Ocean Acoustics Program of ONR and ONR Europe. Only Russian work is support here. Much of the work done under this project is in direct support of work conducted on the influence of bubbles on high-frequency acoustics in shallow water.

REFERENCES

(1) I.P. Smirnov, J.W. Caruthers, and A.I., Khil'ko, Multi-Scale Coherence of the Acoustic Field of a Noise Source in a Random Inhomogeneous Ocean,'in the book: The Forming of Acoustical Waves in an Oceanic Waveguide, Inst. Appl. Phys., RAS, Nizhny Novgorod, (in press), 1997. (2) I.P. Smirnov, A.I. Khil'ko and J.W. Caruthers, "The Role of Weakly Divergent Bundles in Generating Regions with High Correlation in Oceanic Waveguides," in the book The Forming of Acoustical Waves in an Oceanic Waveguide, Inst. Appl. Phys. RAS, Nizhny Novgorod, (in press), 1997. (3) A.I. Khil'ko and J.W. Caruthers, OCEAN ACOUSTIC TOMOGRAPHY: A Review with Emphasis on the Russian Approach, a book in draft with submission to be published by the Inst. Appl. Phys., RAS. (4) I.P. Smirnov, A.I. Khil'ko and J.W. Caruthers, "Partially Coherent Acoustic Tomography in the Ocean," Proceedings of the Third European Conference on Underwater Acoustics, Crete, Greece, pp. 839-844, 1996. (5) D.I. Abrosimov, A.I. Khil'ko and J.W. Caruthers, "Method of Partially Coherent Tomography of Meso-scale Inhomogeneities in the Ocean," *Proceedings of the Third European* Conference on Underwater Acoustics, Crete, Greece, pp. 857-61, 1996. (6) I.V. Smirnov, A.I. Khil'ko, P.A. Elmore, and J.W. Caruthers, "Possibilities for Small-Scale Emission Tomography of Bubble Plumes in Shallow Water.,"Proceedings of the High-Frequency Acoustics in Shallow Water, SACLANTCTR, La Spezia, Italy, 1997. (7) D. Rouseff, F.S. Henyey, J.W. Caruthers, and S.J. Stanic, "Tomographic Reconstruction of Shallow-Water Bubble Field Observations in the Scripps Pier Bubble Experiment," to be presented at the 134th Meeting of the Acoust. Society Amer., abst. published in J. Acoust. Soc. Am., v

102, 1997. (8) O. A. Godin, D. Yu. Mikhin, and D. R. Palmer, "Ocean current monitoring in the coastal zone," accepted for publication in Oceanography. (9) O. A. Godin, "Acoustic mode reciprocity in fluid/solid systems: Implications on environmental sensitivity and horizontal refraction,'Proc. of the 3rd Intl. Conf. on Theor. & Comp. Acoust., Newark, N. J., in July 1997.